

DOCUMENTATION PAGE

Form Approved
OMB No 0704-0188

AD-A210 751

11 1989

1b RESTRICTIVE MARKINGS
NA

3 DISTRIBUTION AVAILABILITY OF REPORT

Distribution unlimited

4 PERFORMING ORGANIZATION REPORT NUMBER(S)

University of California, Los Angeles

5 MONITORING ORGANIZATION REPORT NUMBER(S)

NA

6a NAME OF PERFORMING ORGANIZATION
University of California,
Los Angeles6b OFFICE SYMBOL
(If applicable)

7a NAME OF MONITORING ORGANIZATION

Office of Naval Research

6c ADDRESS (City, State, and ZIP Code)
Department of Biological Chemistry
UCLA School of Medicine
Los Angeles, CA 90024

7b ADDRESS (City, State, and ZIP Code)

800 N. Quincy Street
Arlington, VA 22217-50008a NAME OF FUNDING / SPONSORING
ORGANIZATION
Office of Naval Research8b OFFICE SYMBOL
(If applicable)
ONR

9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER

N00014-86-K-0524

8c ADDRESS (City, State, and ZIP Code)
800 N. Quincy Street
Arlington, VA 22217-5000

10 SOURCE OF FUNDING NUMBER:

PROGRAM
ELEMENT NO
61153NPROJECT
NO
RR04106TASK
NO
441g013WORK UNIT
ACCESSION NO

11 TITLE (Include Security Classification)

Synthetic Sequence Specific Nucleases

12 PERSONAL AUTHOR(S)

David S. Sigman, Ph.D.

13a TYPE OF REPORT
Annual13b TIME COVERED
FROM 8/1/88 TO 7/15/8914 DATE OF REPORT (Year, Month, Day)
7/15/8915 PAGE COUNT
9

16 SUPPLEMENTARY NOTATION

17 COSATI CODES

FIELD	GROUP	SUB-GROUP

18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number)

19 ABSTRACT (Continue on reverse if necessary and identify by block number)

The principal goal of this research was to determine if synthetic peptides derived from the helix-turn-helix domain of DNA binding proteins can be converted into a site specific nuclease by linking it to the nuclease activity of 1,10-phenanthroline-copper.

In this work, we have shown that a synthetic peptide derived from the helix-turn-helix binding domain of the *E. Coli* trp repressor can be converted into a chemical nuclease activity targeted for the aro H operator which is regulated by this control protein. The cutting by the peptide is not as specific as that observed by the chemically derivatized parent protein. Preferred scission is observed because DNA stabilizes the random coil protein in a helix-turn-helix conformation on its surface (eq.1a-c).

20 DISTRIBUTION / AVAILABILITY OF ABSTRACT

☒ UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT ☐ DTIC USERS

21 ABSTRACT SECURITY CLASSIFICATION

U

22a NAME OF RESPONSIBLE INDIVIDUAL
Dr. M. Marron22b TELEPHONE (Include Area Code)
202-696-476022c OFFICE SYMBOL
ONR

peptide-OP (random coil) \longleftrightarrow peptide-OP(h-t-h)* (eq. 1a
 *helix -turn-helix conformation

peptide-OP(h-t-h) + DNA operator sequence \longleftrightarrow (eq. 1b

peptide-OP (h-t-h)-- Operator sequence

Sequence specific scission would result from the following reaction:

peptide-OP (h-t-h)--Operator sequence \rightarrow (eq. 1c

nicked Operator sequence

Background scission is observed because of the high concentration of the peptide existing as a random coil

peptide-OP (random coil) + random sequence DNA \rightarrow
 nicked DNA products (eq. 2

While our work suggests that 1,10-phenanthroline-modified peptides may not prove to be useful as reagent, it demonstrates that DNA binding can stabilize a 21 amino acid peptide in the helix-turn-helix conformation. It has also demonstrated that intact DNA binding proteins can be chemically be converted to specific stable nucleases.

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification _____	
By _____	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	



AMZEL, L. Mario
Department of Biophysics
Johns Hopkins School of Medicine
725 North Wolfe Street
Baltimore, MD 21205

ANDERSEN, Niels H.
Department of Chemistry
University of Washington
Seattle, WA 98195

ARNOLD, Frances H.
Dept of Chemical Engineering
California Institute of Technology
Pasadena, CA 91125

AUGUST, J. Thomas
Department of Pharmacology
Johns Hopkins Medical School
725 North Wolfe Street
Baltimore, MD 21205

BEVERIDGE, David L.
Department of Chemistry
Wesleyan University
Hall-Altwater Laboratories
Middletown, CT 06457

BRAMSON, H. Neal
Department of Biochemistry
Univ of Rochester Medical Center
601 Elmwood Avenue
Rochester, NY 14642

BRUCE, Thomas C.
Department of Chemistry
University of California-Santa
Barbara
Santa Barbara, CA 93106

CASE, Steven T.
Department of Biochemistry
Univ of Mississippi Medical Center
2500 North State Street
Jackson, MS 39216-4505

CHANG, Eddie L.
Bio/Molecular Engineering
Naval Research Laboratory
Code 6190
Washington, D.C. 20375-5000

CHRISTIANSON, David W.
Department of Chemistry
University of Pennsylvania
231 South 34th Street
Philadelphia, PA 19104-6323

CORDINGLEY, John S.
Department of Molecular Biology
University of Wyoming
Box 3944 University Station
Laramie, WY 82071

DeGRADO, William F.
E. I. du Pont de Nemours & Co
Central R & D, Experimental Station
P. O. Box 80328
Wilmington, DE 19880-0328

EVANS, David R.
Department of Biochemistry
Wayne State Univ School of Medicine
540 E. Canfield Street
Detroit, Michigan 48201

FEIGON, Juli F.
Department of Chem & Biochemistry
UCLA
405 Hilgard Avenue
Los Angeles, CA 90024-1569

FICHT, Allison R.
Dept of Med Biochem & Genetics
Texas A&M University
College Station, TX 77843

FRAUENFELDER, Hans
Department of Physics
University of Illinois
Urbana, IL 61801

GABER, Bruce
Naval Research Laboratory
Bio/Molecular Engineering Branch
Code 6190
Washington, DC 20375

GETZOFF, Elizabeth D.
Scripps Clinic & Research Foundation
Department of Molecular Biology
10666 North Torrey Pines Road
La Jolla, CA 92037

GOODMAN, Eugene M.
Biomedical Research Institute
University of Wisconsin
P. O. Box 2000
Kenosha, WI 53141

HO, Pui Shing
Department of Biochemistry and
Biophysics
Oregon State University
Corvallis, OR 97331

HOGAN, Michael E.
Baylor Center for Biotechnology
4000 Research Forest Drive
The Woodlands, TX 77381

HONIG, Barry
Columbia University
Dept of Biochem and Molec Bioph
630 West 168th St.
New York, NY 10032

HOPKINS, Paul B.
Department of Chemistry
University of Washington
Seattle, WA 98195

KAHNE, Daniel
Department of Chemistry
Princeton University
Princeton, NJ 08544

KEMP, Robert G.
Chicago Medical School
Dept of Biological Chemistry
3333 Green Bay Rd.
North Chicago, IL 60064

KHORANA, Gobind H.
Department of Biology
MIT
77 Massachusetts Ave.
Cambridge, MA 02139

KIM, Sangtae
Chemical Engineering
University of Wisconsin
1415 Johnson Drive
Madison, WI 53706

LANSBURY, Peter T.
Department of Chemistry
MIT
Cambridge, MA 02139

LAURSEN, Richard A.
Chemistry Department
Boston University
590 Commonwealth Avenue
Boston, MA 02215

LENZ, Robert W.
Chemical Engineering Department
University of Massachusetts
Amherst, MA 01003

LEWIS, Randolph V.
Molecular Biology Department
University of Wyoming
University Station Box 3944
Laramie, WY 82071

LINDSAY, Stuart M.
Department of Physics
Arizona State University
Temp, AZ 85278

LOEB, George I.
David W. Taylor Research Center
Code 2841
Annapolis, MD 21402-5067

MASILAMANI, Divakar
Biotechnology Department
Allied-Signal Inc.
P. O. Box 1021R
Morristown, NJ 07960

McCONNELL, Harden M.
Stanford University
Department of Chemistry
Stanford, CA 94305

McELROY, William D.
Department of Chemistry
University of California - San Diego
La Jolla, CA 92093-0601

MERTES, Kristin Bowman
University of Kansas
Dept of Chemistry
Lawrence, Kansas 66045

NAGUMO, Mark
Bio/Molecular Engineering Branch
Naval Research Laboratory
Code 6190
Washington, DC 20375-5000

OLIVERA, Baldomero M.
Department of Biology
University of Utah
Salt Lake City, UT 84112

PABO, Carl O.
Department of Biophysics
Johns Hopkins University
School of Medicine
Baltimore, MD 21205

PRENDERGAST, Franklyn G.
Dept of Biochemistry & Molec Biol
Mayo Foundation
200 First St. SW
Rochester, MN 55905

PUGH, Jr., Edward N.
Department of Psychology
University of Pennsylvania
3815 Walnut Street
Philadelphia, PA 19104-6196

RACKOVSKY, Shalom R.
Department of Biophysics
University of Rochester
School of Medicine and Dentistry
Rochester, NY 14642

RAJAN, K. S.
Illinois Institute of Technology
Research Institute
10 W. 35th St.
Chicago, IL 60616

REINISCH, Lou
Laser Biophysics Center
Uniformed Services University
4301 Jones Bridge Road
Bethesda, MD 20814

RICH, Alexander
MIT Department of Biology
Cambridge, MA 02139

RICHARDS, J. H.
California Institute of Technology
Division of Chemistry and Chemical
Engineering
Pasadena, CA 91125

ROTHSCHILD, Kenneth J.
Department of Physics
Boston University
590 Commonwealth Avenue
Boston, MA 02215

SCHULTZ, Peter G.
Department of Chemistry
University of California-Berkeley
Berkeley, CA 94720

SEEMAN, Nadrian
Department of Chemistry
New York University
New York, NY 10003

SELSTED, Michael E.
UCLA
Dept of Medicine
37-055 CHS
Los Angeles, CA 90024

SIGMAN, David S.
UCLA School of Medicine
Dept of Biological Chemistry
Los Angeles, CA 90024
SIKES, Steven C.
Department of Biological Sciences
University of South Alabama
Mobile, AL 36688

SINSKEY, Anthony J.
Laboratory of Applied Microbiology
MIT Department of Biology
Cambridge, MA 02139

STEWART, James M.
Department of Chemistry
University of Maryland
College Park, MD 20742

STEWART, John M.
Department of Biochemistry
University of Colorado
Health Science Center
Denver, CO 80262

TURNER, Douglas H.
Department of Chemistry
University of Rochester
Rochester, NY 14627

URRY, Dan W.
Laboratory of Molecular Biophysics
University of Alabama
P. O. Box 311
Birmingham, AL 35294

WAITE, J. Herbert
College of Marine Studies
University of Delaware
Lewes, DE 19958

WARD, Keith B.
Naval Research Laboratory
Code 6030
Washington, DC 20375

WARSHEL, Arieh
Department of Chemistry
University of Southern California
University Park
Los Angeles, CA 90089-0482

WATT, Gerald D.
Dept of Chemistry & Biochemistry
University of Colorado
Campus Box 215
Boulder, CO 80309-0215

FINAL REPORT
SYNTHETIC SEQUENCE SPECIFIC NUCLEASES

ONR N00014-86-K-0524

DAVID S. SIGMAN, PH.D.

**DEPARTMENT OF BIOLOGICAL CHEMISTRY
SCHOOL OF MEDICINE
AND
MOLECULAR BIOLOGY INSTITUTE
UNIVERSITY OF CALIFORNIA
LOS ANGELES, CA 90024**

JULY 14, 1989

The objectives of our initial proposal included:

1) Targeting DNA with native repressor proteins modified with 1,10-Phenanthroline-Copper.

2) Sequence Independent Scission using Peptides as Carriers of 1,10-Phenanthroline- Copper.

3) Synthesis of Sequence Specific Peptides modified with 1,10-Phenanthroline-Copper

1) Targeting DNA with native repressor proteins modified with 1,10-Phenanthroline-Copper.

Experiments with the *E. Coli trp* repressor demonstrated the feasibility of using a protein or a peptide as a carrier of the nuclease activity of 1,10-phenanthroline-copper. (C.-h. B. Chen and D.S. Sigman "Chemical Conversion of a DNA-Binding Protein into a Site-Specific Nuclease," *Science* 237 1197-1201 (1987)). The nuclease activity of the derivatized protein mirrored the binding specificity of the parent protein. Scission was observed at the two operators tested which are regulated by the native protein. In each case, L-tryptophan was required. This amino acid is a corepressor and must bind to the protein in order to achieve site-specific binding. An additional important feature of the cutting activity, essential for its projected use in chromosomal mapping, is that the reagent accomplished double strand scission.

Since one of the central goals of developing semisynthetic restriction endonuclease is to provide new reagents for chromosomal mapping, we are investigating the reactivity of the modified *trp* repressor within an agarose gel matrix. All manipulations of chromosomal size DNA must be carried out in the gel matrix to avoid shearing of the high molecular weight DNA. Presently, we are investigating the scission of the circular *E. coli* genome. Since there are three binding sites for the *trp* repressor in the *E. coli* genome, it should be possible to identify three distinct segments of DNA by pulsed field gel techniques with appropriate probes in Southern blots.

To improve the efficiency of this scission reagent, 1,10-phenanthroline derivatives with long linker arms have been synthesized. They are currently being used to modify the *E. coli trp* repressor in order to improve the efficiency of the reaction.

2) Sequence Independent Scission using Peptides as Carriers of 1,10-Phenanthroline- Copper.

As initially proposed, we sought to examine the reactivity of 1,10-phenanthroline linked to a peptide which assumed a helical structure upon binding to DNA. This proposed line of experimentation led us to investigate the broader question of the influence of substituents of 1,10-phenanthroline on the specificity of the nuclease activity. The following results have been obtained. 2-Substitution blocks scission since the coordination complex cannot undergo the oxidative cycle necessary for reaction. Substituents at the 5-position are readily tolerated except if they are anionic. Substitution at the equivalent 4 and 7 positions with a methyl group yields an active complex; with a phenyl group, there is a change in reaction mechanism leading to fundamentally different nucleolytic activity. Substitution at the 3-position blocks scission. Substitution at the 5-position is not only tolerated but it also does not alter the intrinsic reactivity of the 1,10-phenanthroline-copper for a given DNA sequence if the substituents lack defined specificity themselves. 5-Substituents on 1,10-phenanthroline which have little influence on the specificity of the scission reaction include methyl, phenyl bromo, and acetamido, aminoethyl and amino hexyl groups. The reason for their lack of effect is apparent from the structure of the essential reactive intermediate formed between the tetrahedral 2:1 5-phenyl-1,10-phenanthroline-cuprous complex and DNA presented in Figure 1. In this model of the essential reactive intermediate, the 5-phenyl substituent does not interact directly with the walls of the minor groove. An important consequence of these studies is that the targeted nucleolytic agents should be synthesized by linking the affinity ligand to the 5-position of the phenanthroline.



Figure 1

The phenanthroline derivatives with 5-substituted cationic groups exhibited similar sequence preferences as the neutral derivatives but generally cut the less reactive sites more efficiently. A more even digestion pattern is consistent with minor groove binding of the coordination complex since the phosphodiester backbone generates a negative potential well. Therefore, cationic phenanthroline derivatives which form a cuprous complexes with net charges between +3 and +4 bind more efficiently than complexes formed with neutral phenanthrolines and (e.g. 5-phenyl-1,10-phenanthroline) with net charge of +1.

In order to examine the reaction of 1,10-phenanthroline-copper linked to a sequence neutral binding peptide, the coordination complex was attached to RecA, a bacterial protein which plays a central role in recombination and binds single and double stranded DNA in a sequence independent manner. Substitution of the phenanthroline at the 5 position with the RecA protein generated a very efficient scission reagent but one which lost the sequence specificity characteristic of the other 1,10-phenanthroline-copper complexes. The tight binding of the Rec A protein must override the small differences in binding affinity which lead to the sequence dependent reactivity. The experiments summarized in this section demonstrate that substitution at the 5-position of phenanthroline does not interfere with the nuclease activity. Moreover, these experiments demonstrated that the linkage of a small protein to the 1,10-phenanthroline-copper complex does not inhibit the nucleolytic activity either by scavenging cupric ion or quenching the oxidative intermediate essential for the DNA scission reaction.

3) Synthesis of sequence specific peptides modified with 1,10-phenanthroline copper.

In contrast to the results summarized above, derivatization of 1,10-phenanthroline at the 5-position with Hoechst dye 33258 yielded a 1,10-phenanthroline derivative with a scission specificity which reflected the A-T specificity of this fluorescent cytological stain (Figure 2). The cutting is particularly efficient because the dye binds within the minor groove.




Figure 2

Hoechst dye 33258 is the first small organic ligand which we have studied that can redirect the cutting activity of 1,10-phenanthroline-copper

The experiments with Hoechst dye, as well as with the RecA protein, indicated a peptide derived from binding domain of a DNA binding protein should be able to target the chemical nuclease activity for site specific cleavage if these peptides had any affinity for DNA. Initially, it was proposed to synthesize peptides derived from the DNA binding domains of phage repressors. However, since we had demonstrated that the *trp* repressor could be transformed into a site specific nuclease, peptides derived from the DNA binding domain of this protein were synthesized instead. In this way, the relative efficiency of cutting of an operator sequence by the intact protein and the peptide could be compared. With the resources available, it was possible to synthesize the following peptides:

Wild type helix-turn-helix peptide

OP-Cys-Gln-Arg-Glu-Leu-Lys-Asn-Glu-Leu-Gly-Ala-Gly-Ile-Ala-Thr-Ile-
Thr-Arg-Gly-Ser-Asn -NH



Chemically mutated helix-turn-helix peptide

OP-Cys-Gln-Arg-Glu-Leu-Lys-Asn-Glu-Leu-Gly-Val-Gly-Ile-Ala-Thr-Ile-
Thr-Arg-Trp-Ser-Asn -NH


These peptide sequences correspond to the helix-turn-helix domain of the *trp* repressor with the following qualifications. The N-terminal cysteine residue is not part of the sequence of the protein but has been included to facilitate derivatization. The underlined residues at 77 and 85 in the wild type repressor are alanine and glycine, respectively. These have been substituted by valine and tryptophan, respectively for the following reasons: a) one mutant *trp* repressor with valine at 77 has a very high affinity for the operator sites; 2) the X-ray structure has shown that L-tryptophan, which must bind to the protein for the repressor to have affinity for DNA, interacts near this glycine and may be essential for stabilizing the conformation of the protein. The peptide derived from the wild-type sequence does not cause sequence dependent scission. Its scission pattern is that of unsubstituted 1,10-phenanthroline-copper.

Each peptide was a) dimerized by forming disulfide bonds at the N-terminal cysteine residue; and b) derivatized by 5-iodoacetyl-1,10-phenanthroline. The interaction of these various synthetic products with the *E. coli* *aroH* operator, one of three regulated by the *E. coli trp* repressor, was studied using DNase footprinting. Under conditions in which the native protein binds with high affinity, none of these peptides showed any sequence specific interaction with the target DNA using the DNase I footprinting assay (Fig 3).

The failure to observe high affinity binding does not necessarily preclude sequence specific cutting by the peptide. The free energy of binding of the peptide for its nucleotide sequence could stabilize the peptide in the conformation competent for DNA binding as indicated in the simple scheme below.



Sequence specific scission would result from the following reaction:



A direct consequence of this simple scheme is that a peptide would bind to a specific sequence more weakly than a protein at comparable concentrations. Since site specific cutting is a reflection of binding, peptide directed cutting would not be expected to be as strong as that of protein directed cutting when presented at equivalent concentrations. The specificity of the cutting (i.e. the site of cutting) would depend on the reactivity of the 1,10-phenanthroline-modified random coil form of the protein free in solution.

In Figure 3, the scission of the nontemplate strand of the *aro* H operator using the OP-Cu derivatized *trp* repressor, underivatized OP-Cu, and OP-Cu derivatized chemically mutated peptide are compared. The products of the scission by the copper complex of OP-linked to the peptide were different from those of the unsubstituted 1,10-phenanthroline-copper exactly in the region in which the derivatized *trp* repressor cuts the nontemplate strand of the *aro* H operator most strongly (Fig 3). This was confirmed by comparing densitometric scans of the digestion pattern of OP-Cu, the peptide linked to OP-Cu and the *trp* repressor derivatized with OP-Cu (Fig.4). The targeted scission by the peptide indicated that the specific DNA sequence is able to stabilize the peptide in a precise conformation.

Peptide-Directed Scission of aro H




Figure 3

Peptide Directed Scission of aro H




Figure 4

Even though enhanced cutting at the operator sites is evident, substantial background cutting is evident. This cutting is due to 1,10-phenanthroline linked to the peptide in the random coil conformation which like other 5-substituted phenanthrolines, reacts similarly to the unsubstituted 1,10-phenanthroline. Since the peptide is present at large excess relative to the target DNA, it is the predominant form of the peptide and 1,10-phenanthroline in the incubation mixture. As a result the large excess of peptide-OP (random coil) will result in the following reaction



nicked DNA products {eq. 2

The conformational instability of this peptide will limit the utility of peptides as sequence specific scission reagents. Binding energies between DNA and the peptide are not large enough to stabilize the conformation of the peptide at equivalent concentrations. *Specific cleavage will be achieved only if the peptide-OP (h-t-h) is intrinsically stable in the helix-turn-helix motif free in solution.* Because of the intrinsic affinity of 1,10-phenanthroline-copper for DNA, it might not be possible to suppress the background reaction of the OP-Cu-peptide. Like the 21 unit described here, the 30 amino acid peptide derived from the DNA binding domain of the transcription factor TF IIIA also fails to exhibit site specific binding for its target sequence using a footprinting assay.

The significant conclusion of our study is that it demonstrates that DNA binding can stabilize a 21 amino acid peptide in the helix-turn-helix conformation.